

Color Evaporation Description of Inelastic Photo-Production of J/ψ at HERA

O. J. P. Éboli,^{1,*} E. M. Gregores,^{2,†} and F. Halzen^{3,‡}

¹*Instituto de Física, Universidade de São Paulo, São Paulo – SP, Brazil.*

²*Instituto de Física Teórica, Universidade Estadual Paulista, São Paulo – SP, Brazil.*

³*Department of Physics, University of Wisconsin, Madison – WI, USA.*

The H1 Collaboration recently reported a new analysis on the inelastic photo-production of J/ψ mesons at DESY HERA ep collider. We show that these new experimental results are well described by the Color Evaporation Model for quarkonium production. Moreover, this new data requires the introduction of resolved photon contributions in order to explain the results on small charmonium energy fraction, indicating that colored $c\bar{c}$ pairs also contribute to the process.

PACS numbers: 13.60.Le, 25.20.Lj

I. INTRODUCTION

The H1 Collaboration recently reported an analysis on the inelastic photo-production of J/ψ mesons [1] where their new data was confronted with the color singlet [2] and color octet [3] models. We compare the H1 data with the color evaporation model (CEM) predictions for quarkonium production showing that it provides a good description of this new set of data. Moreover, this new data probes for the first time the small charmonium energy fraction (z) region, requiring the introduction of resolved photon contributions in order to explain the results. These contributions are basically due to color octet configurations, confirming the correctness of their inclusion in the charmonium production mechanism.

Previous measurements of the inelastic photo-production of charmonium at HERA [4, 5] appeared to have ignited a charmonium crisis. The color singlet model (CSM) to onium production cross section fits the data for large charmonium energy fraction z , where color octet models seemed to fail. This fact is, however, in qualitative disagreement with a wealth of information that exists on charmonium production by other initial states. We suggested [6] that this discrepancy is due to the neglect of non-perturbative effects that are important at large z . Implementing a phenomenological parametrization of these effects in a scheme originally developed for Drell-Yan phenomenology, we illustrated how agreement with data could be achieved. In this work, we employ the procedure of [6] to mimic the non-perturbative effects.

II. THEORETICAL FRAMEWORK

It is clear nowadays that J/ψ production is a two-step process where a heavy quark pair is produced first, fol-

lowed by the non-perturbative formation of the colorless asymptotic state. The predictions of the once conventional treatment of color in perturbative QCD calculations, *i.e.*, the CSM [7], for the charmonium production at the Fermilab Tevatron is at variance with the available data[8]. As a consequence, color octet as well as singlet $c\bar{c}$ states contribute to the production of J/ψ . Two formalisms have been proposed to incorporate these features: the Non-Relativistic QCD, also known as COM [9], and the renewed CEM scheme [10, 11], also known as Soft Color Interactions (SCI) [12]. The original CEM [13] actually predates the color singlet approach, and had been abandoned for no good reason. Recent measurements of the polarization of bound charm and beauty mesons, seems to disfavor the COM framework [14].

The color evaporation model simply states that charmonium production is described by the same dynamics as $D\bar{D}$ production, *i.e.*, by the formation of a colored $c\bar{c}$ pair. Rather than imposing that the $c\bar{c}$ pair is in a color-singlet state in the short-distance perturbative diagrams, it is argued that the appearance of color-singlet asymptotic states solely depends on the outcome of large-distance fluctuations of quarks and gluons. These large-distance fluctuations are probably complex enough for the occupation of different color states to approximately respect statistical counting. In other words, the formation of color-singlet states is a non-perturbative phenomenon. In fact, it does not seem logical to enforce the color-singlet property of the $c\bar{c}$ pair at short distances, given that there is an infinite time for soft gluons to readjust the color of the pair before it appears as an asymptotic ψ , χ_c or, alternatively, $D\bar{D}$ state. It is indeed hard to imagine that a color-singlet state formed at a range m_ψ^{-1} automatically survives to form a ψ . This approach to color is also used to formulate a successful prescription for the production of rapidity gaps between jets at Tevatron [17, 18, 19, 20, 21] and HERA [20, 21, 22, 23].

Although far more restrictive than other proposals, the CEM successfully accommodates all features of charmonium production [3, 15, 16]. It predicts that the sum of the cross section of all onium and open charm states is

*Electronic address: eboli@fma.if.usp.br

†Electronic address: gregores@ift.unesp.br

‡Electronic address: halzen@pheno.physics.wisc.edu

described by [10, 17]

$$\sigma_{\text{onium}} = \frac{1}{9} \int_{2m_c}^{2m_D} dM_{c\bar{c}} \frac{d\sigma_{c\bar{c}}}{dM_{c\bar{c}}}, \quad (1)$$

and

$$\sigma_{\text{open}} = \frac{8}{9} \int_{2m_c}^{2m_D} dM_{c\bar{c}} \frac{d\sigma_{c\bar{c}}}{dM_{c\bar{c}}} + \int_{2m_D} dM_{c\bar{c}} \frac{d\sigma_{c\bar{c}}}{dM_{c\bar{c}}}, \quad (2)$$

where $M_{c\bar{c}}$ is the invariant mass of the $c\bar{c}$ pair. The factor $1/9$ stands for the probability that a pair of charm quarks formed at a typical time scale $1/M_\psi$ ends up as a color singlet state after exchanging an uncountable number of soft gluons with the reaction remnants. One attractive feature of this model is the relation between the production of charmonium and open charm which allows us to use the open charm data to normalize the perturbative QCD calculation, and consequently, to make more accurate predictions for charmonium cross sections.

The fraction ρ_ψ of produced onium states that materialize as ψ ,

$$\sigma_\psi = \rho_\psi \sigma_{\text{onium}}, \quad (3)$$

has been inferred from low energy measurements to be a constant [24, 25]. From the charmonium photo-production, we determined that $\rho_\psi = 0.43\text{--}0.5$ [11]; a value that can be accounted for by statistical counting of final states [12]. The fact that all ψ production data are described in terms of this single parameter, fixed by J/ψ photo-production, leads to parameter free predictions for Z -boson decay rate into prompt ψ [26], and to charmonium production cross section at Tevatron [27] and HERA [6], as well as in neutrino initiated reactions [28]. These predictions are in agreement with the available data.

III. INELASTIC PHOTO-PRODUCTION OF CHARMONIUM

Application of the CEM scheme to inelastic charmonium photo-production is straightforward. According to the parton model, the cross section for the J/ψ photo-production at a given center-of-mass energy W is

$$\sigma_{\gamma p \rightarrow J/\psi X}(W) = \int \int f_{A/\gamma}(x_A) f_{B/p}(x_B) \hat{\sigma}_{AB \rightarrow J/\psi X}(\hat{s}) dx_A dx_B, \quad (4)$$

where the subprocess cross section $\hat{\sigma}$ is given by Eqs. (1) and (3). Here, $\sqrt{\hat{s}} = \sqrt{x_A x_B} W$ is the center-of-mass energy of the subprocess $AB \rightarrow J/\psi X$, and $f_{A/\gamma}$ ($f_{B/p}$) is the distribution function of the parton A (B) in the photon (proton). For direct photon interactions ($A = \gamma$) we have $f_{A/\gamma}(x_A) = \delta(x_A - 1)$.

An important kinematical variable is

$$z \equiv \frac{P_{J/\psi} \cdot P_p}{P_\gamma \cdot P_p}, \quad (5)$$

where $P_{J/\psi, \gamma, p}$ is the four-momentum of the J/ψ , photon, or proton, respectively. In the proton rest frame, z is the fraction of photon energy carried by the J/ψ .

The lowest order CEM contribution for the charmonium production is the direct photon process $\gamma g \rightarrow c\bar{c}$. However, it is important only for $z \approx 1$. For the range of z we are interested, the direct photon contribution is dominated by the diagrams depicted in the Fig. 1(a). The charm quark pair in γg fusion can be produced in both color singlet and octet configurations, while γq fusion leads only to colored $c\bar{c}$ pairs. Besides the direct photon–gluon and photon–quark reactions, we also included resolved photon processes, which proceed via quark–quark, quark–gluon, and gluon–gluon fusion into $c\bar{c}$ +quark (gluon), as exemplified in Fig. 1(b).




FIG. 1: Processes contributing to direct (a) and resolved (b) charmonium photo-production.

The higher order processes, like the ones in Fig. 1, have to be evaluated with some caution in the region of rela-

tively small J/ψ transverse momentum and large z . For small t -channel momentum transfer ($\hat{t} \rightarrow 0$), the gluon exchange diagrams in Fig. 1 represent the QCD evolution of the initial state gluon distribution functions, and not higher order gluon exchange diagrams. Therefore, the contribution of the gluon-exchange diagrams where $\hat{t}$ is soft, has already been taken into account by the leading order $\gamma g \rightarrow c \bar{c}$ diagram [29]. By itself, when evaluated at tree level, the diagrams in Fig. 1 lead to a divergence for $z \rightarrow 1$, that appears as an unphysical growth of the cross section for $z \lesssim 1$. Although the double counting is relevant only for vanishing p_T and $z = 1$, the enhancement of the cross section associated with the collinear divergence will, in practice, affect the calculation until p_T and z -values become truly perturbative.

Presently, complete higher order QCD calculations are not available for the processes we are interested in. Therefore, we phenomenologically took into account non-perturbative QCD corrections in large z charmonium photo-production by introducing the following parametrization to mimic the non-perturbative contributions [6]:

$$\frac{d^2\sigma}{dp_T dz} = [1 - F(Q_0, p_T)][1 - G(Q_0, z)] \frac{d^2\sigma_{tree}}{dp_T dz} \quad (6)$$

with

$$F(Q_0, p_T) = e^{-\frac{p_T^2}{k_T^2}} \quad (7)$$

and

$$G(Q_0, z) = e^{-\frac{1-z}{z_0 z}}, \quad (8)$$

where σ_{tree} is the tree level perturbative cross section, $k_T^2 = z(Q_0^2 + 4m_c^2) - 4m_c^2$, and $(1 - z_0) = (p_T^2 + 4m_c^2)/(Q_0^2 + 4m_c^2)$ are positive definite or null. The parameter Q_0 indicates the value of the momentum transfer $\hat{t}$ where the higher order diagrams contribute in the truly perturbative regime [30]. We found in [6] that $Q_0 = 2m_c$ best describes the data, and it is this value that we used in the present work.

IV. RESULTS

In this work we evaluated numerically the tree level scattering amplitudes using the Madgraph [31] and Helas [32] packages. The phase space integration was performed using the adaptative Monte Carlo program VEGAS [33].

There is a wide choice of QCD parameters appearing in the evaluation of charmonium photo-production, which leads to a large theoretical uncertainty. For the sake of definiteness, we present only the uncertainty that arises from possible choices of the charm quark mass, keeping all others QCD parameters fixed. All our results are presented as a central value surrounded by an error band corresponding to the choice of the charm quark mass as

$m_c = 1.3 \pm 0.1$ GeV. We used the GRV-94 LO [34] parameterization of the proton structure functions and GRV-G LO [35] for the photon parton density. For both structure functions, we set the factorization scale as $\mu_F = \sqrt{\hat{s}}$. We evaluated the running of the strong coupling constant in leading order with four active flavors using $\Lambda_{QCD} = 300$ MeV, renormalization scale $\mu_R = \sqrt{2}m_c$ for direct processes, and $\mu_R = \sqrt{\hat{s}}$ for resolved processes. We also used $\rho_\psi = 0.5$ as the J/ψ fraction of all charm bound states.

The H1 Collaboration performed their analysis using data in which $Q^2 < 1$ GeV², and subdivided their data into several different kinematical regions, in order to better determine the region where perturbative QCD calculations furnish a reliable description of the data.




FIG. 2: Total cross section as function of $W_{\gamma p}$ for $p_T > 1$ GeV: (a) $0.3 < z < 0.9$ and (b) $0.3 < z < 0.8$. The shaded band shows the theoretical prediction for $m_c = 1.3 \pm 0.1$ GeV.

We collated the CEM predictions with the H1 results using the cuts applied to the experimental data. Initially we analyzed the behavior of the total cross section as function of the γp center-of-mass energy (W) for two different z regions, requiring a minimum J/ψ transverse momentum $p_T > 1$ GeV. In Fig. 2(a) we show the CEM predictions for $0.3 < z < 0.9$, and in Fig. 2(b) we present our results for $0.3 < z < 0.8$. As we can see from these figures, moving the upper limit in z alone does not change much the shape of the curve, and within theoretical uncertainties, agreement is found for both regions.

We display in Fig. 3, the CEM predictions for the z spectrum, requiring $p_T > 1$ GeV and a reaction center-of-mass energy in the range $120 < W_{\gamma p} < 260$ GeV. In this figure, the experimental points represented by triangles and squares stand for two different datasets; see Ref. [1] for details. The direct photon contribution is represented by the dashed line, and the resolved one by the dotted line; the solid line displays the sum of direct and resolved contributions. The shaded area corresponds to the total CEM prediction for the charm quark mass described above. As we can see from this figure, the agreement of the CEM with data is quite good. Moreover, the data at low and medium z do require the introduction of




FIG. 3: Differential cross section as function of the inelasticity parameter z for $p_T^2 > 1 \text{ GeV}^2$ and $120 < W_{\gamma p} < 260 \text{ GeV}$. The dashed line stands for the contribution from direct processes while the dotted line shows the contribution from resolved processes. The solid line is the sum of both contributions. The shaded band shows the theoretical uncertainty on the value of the charm mass ($m_c = 1.3 \pm 0.1 \text{ GeV}$).

resolved processes in order to explain the results. This is a clear signal that colored charm quark pairs contribute to the J/ψ production.

In Fig. 4 we present the p_T^2 distribution for the low z sample $0.05 < z < 0.45$ which corresponds to the triangles in Fig. 3. For these values of z we expect the theoretical uncertainties due to higher order corrections to be small, and we can see that the CEM and data agree well. Consequently, the disagreement between data and theory at large z is indeed due to the importance of higher order non-perturbative corrections.




FIG. 4: Squared transverse momentum p_T^2 distribution for data collected in the very inelastic region $0.05 < z < 0.45$ and $120 < W_{\gamma p} < 260 \text{ GeV}$. The shaded band shows the theoretical uncertainty on the value of the charm mass ($m_c = 1.3 \pm 0.1 \text{ GeV}$).

Higher order effects are sizeable on the medium and high z regions. Therefore, the H1 Collaboration divided their data sample into several p_T and z regions for a

center-of-mass energies in the range $60 < W_{\gamma p} < 240 \text{ GeV}$. This allows us to better compare data with theory.




FIG. 5: Differential cross section as function of the inelasticity parameter z for $p_T > 1, 2$, and 3 GeV , divided by a factor 1, 10, and 100 respectively. The shaded bands show the theoretical prediction obtaining varying the charm mass ($m_c = 1.3 \pm 0.1 \text{ GeV}$).

In Fig. 5, we compare the CEM predictions for the z spectrum with data for three different values of the J/ψ minimum transverse momentum, *i.e.*, $p_T > 1, 2$, and 3 GeV . The curves have been divided by factors 1, 10, and 100, respectively, in order to help visualization. There is an overall agreement between data and theory, except for the highest bin in z . Moreover, removing our regularization procedure only worsens the theoretical results [6]. Increasing the minimum value of the transverse momentum does not improve the quality of the fitting, what means that our parametrization of the higher order effects has correctly incorporated the dependence on minimum p_T .




FIG. 6: Squared transverse momentum p_T^2 distribution for $0.3 < z < 0.9$ and center-of-mass energy $60 < W_{\gamma p} < 240 \text{ GeV}$. The shaded band shows the theoretical uncertainty on the value of the charm mass ($m_c = 1.3 \pm 0.1 \text{ GeV}$).

Fig. 6 contains the J/ψ squared transverse momentum




FIG. 7: Squared transverse momentum p_T^2 spectrum for $60 < W_{\gamma p} < 240$ and data collected in three different inelastic bins: $0.75 < z < 0.9$ (upper), $0.6 < z < 0.75$ (middle), and $0.3 < z < 0.6$ (lower). To help visualization the curves have been scaled as 1, 1/10, and 1/100 from top to bottom. The shaded bands show the theoretical uncertainty on the value of the charm mass ($m_c = 1.3 \pm 0.1$ GeV).

distribution taking into account the medium and high z data. The agreement between theory and data is quite satisfactory. However, the shape of the spectrum shows some disagreement at the very low p_T bins. In order to understand what is happening, let us consider this distribution for three different z bins, namely $0.75 < z < 0.9$, $0.6 < z < 0.75$, and $0.3 < z < 0.6$; see Fig. 7 where the curves have also been divided by factors 1, 10, and 100, respectively, to help visualization. We can learn from this last figure that the agreement between CEM predictions

and data improves for low z regions. This fact can be understood as a limitation of the proposed parameterization to completely mimic higher order QCD contributions when we approach the elastic region. It also implies that the observed small discrepancy with data for the highest bin on z must be credited to the lack of a complete QCD calculation and is not related to the Color Evaporation approach to describe quarkonium production.

V. CONCLUSIONS

We showed that the Color Evaporation Model describes the available data on J/ψ photo-production, provide we include high order QCD corrections at high inelasticities z . Moreover, the newly available data at low z provides a clear proof of the importance of colored $c\bar{c}$ pairs to the production of charmonium, since the data on this region can only be explained considering resolved photon processes, which lead to colored $c\bar{c}$ pairs.

Acknowledgments

This research was supported in part by the University of Wisconsin Research Committee with funds granted by the Wisconsin Alumni Research Foundation, by the U.S. Department of Energy under grant DE-FG02-95ER40896, by Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP), by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), and by Programa de Apoio a Núcleos de Excelência (PRONEX).

-
- [1] C. Adloff *et al.* [H1 Collaboration], DESY-02-059, May 2002 [hep-ex/0205064].
 - [2] M. Kramer, Nucl. Phys. B **459**, 3 (1996).
 - [3] M. Kramer, Prog. Part. Nucl. Phys. **47**, 141 (2001).
 - [4] S. Aid *et al.* [H1 Collaboration], Nucl. Phys. B **472**, 3 (1996).
 - [5] J. Breitweg *et al.* [ZEUS Collaboration], Z. Phys. C **76**, 599 (1997).
 - [6] O. J. Eboli, E. M. Gregores, and F. Halzen, Phys. Lett. B **451**, 241 (1999).
 - [7] R. Baier and R. Ruckl, Z. Phys. C **19**, 251 (1983).
 - [8] F. Abe *et al.* [CDF Collaboration], Phys. Rev. Lett. **69**, 3704 (1992); *ibid* **79**, 572 (1997); *ibid* **79**, 578 (1997); S. Abachi *et al.* [DØ Collaboration], Phys. Lett. B **370**, 239 (1996).
 - [9] G. T. Bodwin, E. Braaten, and G. P. Lepage, Phys. Rev. D **51**, 1125 (1995) [Erratum-*ibid.* D **55**, 5853 (1997)].
 - [10] J. F. Amundson, O. J. Eboli, E. M. Gregores, and F. Halzen, Phys. Lett. B **372**, 127 (1996).
 - [11] J. F. Amundson, O. J. Eboli, E. M. Gregores, and F. Halzen, Phys. Lett. B **390**, 323 (1997).
 - [12] A. Edin, G. Ingelman, and J. Rathsmann, Phys. Rev. D **56**, 7317 (1997).
 - [13] H. Fritzsch, Phys. Lett. B **67**, 217 (1977); F. Halzen, Phys. Lett. B **69**, 105 (1977); F. Halzen and S. Matsuda, Phys. Rev. D **17**, 1344 (1978).
 - [14] T. Affolder *et al.* [CDF Collaboration], Phys. Rev. Lett. **85**, 2886 (2000).
 - [15] C. B. Mariotto, M. B. Gay Ducati, and G. Ingelman, Eur. Phys. J. C **23**, 527 (2002).
 - [16] G. A. Schuler and R. Vogt, Phys. Lett. B **387**, 181 (1996).
 - [17] O. J. Eboli, E. M. Gregores, and F. Halzen, *Proceedings of the 26th International Symposium on Multiparticle Dynamics (ISMD 96)*, Faro, Portugal, 1996.
 - [18] O. J. Eboli, E. M. Gregores, and F. Halzen, Nucl. Phys. Proc. Suppl. **71**, 349 (1999).
 - [19] O. J. Eboli, E. M. Gregores, and F. Halzen, Phys. Rev. D **58**, 114005 (1998).
 - [20] O. J. Eboli, E. M. Gregores, and F. Halzen, Phys. Rev. D **61**, 034003 (2000).
 - [21] O. J. Eboli, E. M. Gregores, and F. Halzen, Nucl. Phys. Proc. Suppl. **99A**, 257 (2001).
 - [22] W. Buchmüller, Phys. Lett. B **353**, 335 (1995); W. Buchmüller and A. Hebecker, Phys. Lett. B **355**, 573 (1995).
 - [23] A. Edin, G. Ingelman, and J. Rathsmann, Phys. Lett. B **366**, 371 (1996).
 - [24] R. Gavai *et al.*, Int. J. Mod. Phys. A **10**, 3043 (1995).

- [25] G. Schuler, report CERN-TH.7170/94.
- [26] O. J. Eboli, E. M. Gregores, and F. Halzen, Phys. Lett. B **395**, 113 (1997).
- [27] O. J. Eboli, E. M. Gregores, and F. Halzen, Phys. Rev. D **60**, 117501 (1999).
- [28] O. J. Eboli, E. M. Gregores, and F. Halzen, Phys. Rev. D **64**, 093015 (2001).
- [29] P. Nason, S. Dawson, and R. K. Ellis, Nucl. Phys. B **303**, 607 (1988).
- [30] E. Berger and D. Jones, Phys. Rev. D **23**, 1521 (1981).
- [31] W. Long and T. Steltzer, Comput. Phys. Commun. **81**, 357 (1994).
- [32] H. Murayama, I. Watanabe, and K. Hagiwara, KEK report 91-11.
- [33] G. P. Lepage, CLNS-80/447.
- [34] M. Gluck, E. Reya, and A. Vogt, Z. Phys. C **67**, 433 (1995).
- [35] M. Gluck, E. Reya, and M. Stratmann, Phys. Rev. D **51**, 3220 (1995).